

Andrew Wymore, Travis Landon, Lydia Brown and Dr. Chris Walser

Department of Biology, The College of Idaho, 2112 Cleveland Blvd, Caldwell, ID, 83605

## Introduction:

The Columbia River redband trout (*Oncorhynchus mykiss gairdneri*) is native to the Columbia River Basin east of the Cascades and below barrier falls (e.g. Shoshone Falls, Idaho) (Behnke, 1992). Like many salmonids, Columbia redband trout are threatened by introgression with hatchery stocks. Dry Creek (Figure 1), however, is the only stream along the Boise front known to support genetically pure redband trout (Richins, Walsh, and Walser, in preparation).


Dry Creek is an IRES (intermittent river and ephemeral stream) system, and as such, exhibits large seasonal fluctuations in stream flow. IRES systems are the most common water bodies in many parts of the world (Datry et al. 2017). Interestingly, far less research has been done on these systems than perennial waterways. Furthermore, climate change scenarios for semi-arid western North America predict longer and drier summers with decreases in mean annual stream flow (Seager et al. 2013). Changes in stream physicochemistry resulting from flow decline can be extremely stressful to aquatic organisms (Datry et al. 2017).

## Research Objective:


To investigate the impact of flow cessation on stream physicochemistry, macroinvertebrate drift, and Columbia River redband trout (*Oncorhynchus mykiss gairdneri*) condition.

## Materials and Methods:


- 10 reaches of Dry Creek were investigated from June–September 2019. (Figure 1)
- Current velocity was measured weekly at 60% of depth using a Swiffer Model 2100 current velocity meter.
- HOBO U26 dissolved oxygen (DO) and HOBO Pendant temperature loggers were deployed in the deepest pool in each study reach.
- Macroinvertebrates were collected weekly with a drift-net.
- Fish were collected using a Smith-Root Model 12 backpack electrofisher on three separate occasions for each study reach.
- Total length (TL-mm) and weight (g) of each fish was measured with a fish measuring board and OHAUS model SPX8200 scale, respectively, during each sampling event.
- Fish were PIT tagged upon initial capture.
- Condition Factor (K) was determined for each fish using the Fulton equation (Froese, 2006).
- Statistical analyses were performed using SigmaPlot v13.0.




**Figure 1.** Location of Dry Creek study sites. Dry Creek experiences drying every summer—beginning in the downstream reaches. The headwaters of Dry Creek remain perennial. Study sites span a distance of about 5-km.




**Figure 2.** Dry Creek current velocity (m/s) at sites 1, 6, and 9 from June through August 2019.




**Figure 3a.** Site 1 dissolved oxygen and temperature profile.




**Figure 3b.** Site 6 dissolved oxygen and temperature profile.




**Figure 3c.** Site 9 dissolved oxygen and temperature profile.



**Figure 4.** Estimated redband trout abundance across space and time in Dry Creek for three sample events per site.



**Figure 5.** Changes in mean redband trout condition (K) in Dry Creek over the study period. \* $p < 0.05$



**Figure 6.** Macroinvertebrate drift density over the study period at sites 1, 6, and 9, collected weekly.



Flow cessation in Dry Creek. These photographs were taken at the same location in June (left) and August (right).

## Study Conclusions:

- Current velocity decreased across all sites during the study period with Site 1 experiencing complete flow cessation. (Figure 2)
- Mean daily water temperatures increased during the summer across all sites but were within an acceptable range for redband trout. (Figure 3)
- The most downstream pool experienced the lowest dissolved oxygen (3 mg/l) upon flow cessation. (Figure 3a)
- Redband trout were able to persist in pools with dissolved oxygen levels below 6 mg/l for at least 9 days. (Figures 3a and 3b)
- Redband trout were more abundant in upstream than downstream reaches. (Figure 4)
- Redband trout captured early in the study were in significantly better overall health than those caught later. (Figure 5)
- Macroinvertebrate drift declined across all sites over time. (Figure 6)
- We attribute the drop in redband trout condition over the summer, in part, to decreased oxygen levels and reduced food availability (macroinvertebrate drift) due to channel constriction and flow reduction.

## Acknowledgments:

This research was made possible by the M.J. Murdock Charitable Trust under grant number SR-201811723

A portion of this research was also supported by the NSF Idaho EPSCoR Program and by the National Science Foundation under award OIA-1757324.

We would also like to thank The College of Idaho Biology Department for providing funds in support of this research.

Finally, we would like to thank Ceanna Schwartz for creating our map.

## References:

- Behnke, R. J. 1992. Native trout of western North America. 275 pp.
- Datry, T., Núria, B., & A. Boulton. 2017. Intermittent Rivers and Ephemeral Streams: Ecology and Management.
- Froese R. 2006. Cube law, condition factor and weight–length relationships: history, meta-analysis and recommendations. *Journal of Applied Ichthyology*, 22:241–253.
- Gamperl, A. K., Rodnick, K. J., Faust, H. A., Venn, E. C., Bennett, M. T., Crawshaw, L. I., Keeley, E. R., Powell, M. S., and H.W Li. 2002. Metabolism, swimming performance, and tissue biochemistry of high desert redband trout (*Oncorhynchus mykiss* ssp.): Evidence for phenotypic differences in physiological function. *Physiological and Biochemical Zoology*, 75(5), 413–431.
- Seager, R., Ting, M. Li, C., Naik, N., Cook, B., Nakamura, J., et al., 2013. Projections of declining surface-water availability for the southwestern United States. *Nature Climate Change* 3, 482–486.



Columbia River redband trout in Dry Creek.